



PATIENT INFORMATION (Complete or Fax Existing Chart) PRESCRIBER INFORMATION

Name: \_\_\_\_\_ DOB: \_\_\_\_\_
Address: \_\_\_\_\_
City, State, Zip: \_\_\_\_\_
Phone: \_\_\_\_\_ Alt. Phone: \_\_\_\_\_
Email: \_\_\_\_\_ SS#: \_\_\_\_\_
Gender:  M  F Weight: \_\_\_\_\_ (lbs) Ht: \_\_\_\_\_
Allergies: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_
State License: \_\_\_\_\_
NPI #: \_\_\_\_\_ DEA: \_\_\_\_\_
Address: \_\_\_\_\_
City, State, Zip: \_\_\_\_\_
Phone: \_\_\_\_\_ Fax: \_\_\_\_\_
Office Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)

Primary Insurance: \_\_\_\_\_
Plan #: \_\_\_\_\_
Group #: \_\_\_\_\_
RX Card (PBM): \_\_\_\_\_
BIN: \_\_\_\_\_ PCN: \_\_\_\_\_

Secondary Insurance (If Applicable): \_\_\_\_\_
Plan #: \_\_\_\_\_
Group #: \_\_\_\_\_
RX Card (PBM): \_\_\_\_\_
BIN: \_\_\_\_\_ PCN: \_\_\_\_\_

CLINICAL INFORMATION

Primary ICD-10 (Please Specify Diagnosis): \_\_\_\_\_ Secondary ICD-10 (Please Specify Diagnosis): \_\_\_\_\_
Tertiary ICD-10 (Please Specify Diagnosis): \_\_\_\_\_
Is the patient on iron, folate and/or vitamin B12 therapy?  Yes  No Is the patient on dialysis?  Yes  No
Has patient received any ESA therapy?  Yes  No If yes, how many weeks of ESA therapy has the patient completed? \_\_\_\_\_ weeks
Patient's hemoglobin (Hgb) level: \_\_\_\_\_ g/dL

ARANESP® ORDERS

Prescription type:  New start  Restart  Continued therapy Total Doses Received: \_\_\_\_\_ Date of Last Injection/Infusion: \_\_\_\_\_

Table with 3 columns: Medication, Dose/Frequency, Refills. Includes options for Aranesp (darbepoetin alfa) Single Dose Vials and Single Dose Prefilled Syringe with various dose and frequency options.

Special Instructions: \_\_\_\_\_

Table with 3 columns: Pre-Medication, Dose/Strength, Directions. Lists acetaminophen, diphenhydramine, and methylprednisolone with their respective dosages and directions for use.

INFUSION REACTION ORDERS

Mild reaction protocol:
[ ] Diphenhydramine 25mg IV, one time, for pruritus.
If symptoms worsen, see orders for moderate to severe reactions.
Moderate reaction protocol:

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation.



Please Fax Completed Form To: 888-340-3699

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Please Send a Copy of The Patient's Up-to-Date Clinical Notes

- Acetaminophen 650mg PO, one time, for pyrexia or rigors
  - Diphenhydramine 50mg IV, one time, for pruritus or urticaria
  - Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms
- If symptoms worsen, see interventions for severe reactions*
- Severe reaction protocol: (Call 911 if initiated):**
- Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)
  - Diphenhydramine 50mg IV, one time, for respiratory symptoms, edema, or anaphylaxis
  - Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis
  - Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms
  - Epinephrine 0.3mg/0.3mL IM into mid-antrolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or worsen

**FLUSHING & LOCKING ORDERS**

Flushing Protocol (>66lbs/33kg)

<b>PIV and Midline:</b> <input checked="" type="checkbox"/> 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion	<b>Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:</b> <input checked="" type="checkbox"/> 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw
---	--

Locking Protocol (>66lbs/33kg)

<b>PIV and Midline:</b> <input checked="" type="checkbox"/> Heparin Sodium 10 units/mL 1mL IV final flush post normal saline flush	<b>PICC:</b> <input checked="" type="checkbox"/> Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush	<b>Implanted Port, Tunneled Catheter, and Non-tunneled Catheter:</b> <input checked="" type="checkbox"/> Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush
---	--	---

**\*\* May substitute Dextrose 5% in Water, or alternative, for 0.9% Sodium Chloride, when indicated due to incompatibility with medications being infused**

**SIGNATURE**

We hereby authorize Infuse IQ LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.

X \_\_\_\_\_  
Prescriber Signature

Date: \_\_\_\_\_

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.